

Derivation of the Entropy-area Relation for the Extremal Black Hole in Four Dimensions

B. B. Deo*

Department of Physics, Utkal University, Bhubaneswar-751004, India

P. K. Jena†

72, Dharma Vihar, Khandagiri, Bhubaneswar-751030, India

(Dated: July 3, 2009)

The relation between entropy and the area of the event horizon of a quantum black hole in four dimensions is derived. The Reissner-Nordstrom metric for a non-rotating, charged black hole is shown to be modified by addition of a new $\frac{1}{r^3}$ term arising from the strong interaction. The resulting entropy-area relation agrees with the Bekenstein-Hawking relation.

PACS numbers: 04.70.Dy, 04.20.-q, 11.25.-w

Keywords: Black hole, General relativity, Strings

1. INTRODUCTION

It has been a long standing problem to derive the Bekenstein-Hawking law that relates the entropy and area of the event horizon of a black hole, using statistical mechanics in terms of the microstates of the black hole. Classically, the strong gravitational field of a black hole prevents the emission of any radiation from it and the entropy associated with the black hole is zero. However, the scenario is changed when the black hole is treated as a quantum system. Quantum mechanical tunneling through the potential barrier permits the creation of particle pairs from vacuum in the strong gravitational field near the black hole. This is analogous to the electron-positron pair creation, from vacuum, by very strong electric fields. Thus quantum black holes can emit radiation into space. This radiation has the characteristic of black body radiation. So, black holes can be treated as thermodynamic systems with temperature (in geometrized units, $c = G = 1$)

$$T = \frac{\kappa}{2\pi}, \quad (1)$$

where κ is the surface gravity of the black hole. When the mass of a black hole increases due to absorption of nearby stars or due to collision with other black holes, it expands and the area increases. The increase in the mass of the black hole is related to the increase in its area by

$$dM = \frac{\kappa}{8\pi} dA, \quad (2)$$

where M is the mass of the black hole and A is the area of its event horizon. In a thermodynamic system, the increase in entropy and energy are related by

$$dE = T dS, \quad (3)$$

where dE is the increase in energy and dS is the increase in entropy. Equations (1,2,3) suggest a relation between the entropy and area of a black hole as

$$\frac{\kappa}{8\pi} dA = T dS = \frac{\kappa}{2\pi} dS,$$

which leads to the Bekenstein-Hawking relation[1, 2]

*Electronic address: bdeo@iopb.res.in

†Electronic address: prasantajena@yahoo.com

$$S = \frac{A}{4}, \quad (4)$$

between entropy and area of a black hole. Although this relation was obtained using thermodynamic considerations, it was not possible to calculate the entropy of the black hole using statistical mechanics that relates the entropy with the micro-states of the black hole.

For a charged, non-rotating black hole of mass M and electric charge Q , the metric (which will be given in detail later) obtained by Reissner and Nordstrom [3, 4] as solution of Einstein's field equation in a spherically symmetric space is given by

$$g_{tt} = -\left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right), \quad g_{rr} = \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right)^{-1}, \quad g_{\theta\theta} = r^2 \quad \text{and} \quad g_{\phi\phi} = r^2 \sin^2 \theta. \quad (5)$$

So, for the Reissner-Nordstrom metric $g_{\mu\nu}$, the line element $d\tau$ is

$$d\tau^2 = g_{\mu\nu} dx^\mu dx^\nu = -\left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right)^{-1} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2). \quad (6)$$

If r_0 and r_1 are the solutions of the quadratic equation

$$\left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right) = \left(1 - \frac{r_0}{r}\right) \left(1 - \frac{r_1}{r}\right) = 0, \quad (7)$$

so that $r_0 + r_1 = 2M$ and $r_0 r_1 = Q^2$, we have

$$r_{0,1} = M \pm \sqrt{M^2 - Q^2}. \quad (8)$$

The equation(6) can now be recast, in terms of r_0 and r_1 , as

$$d\tau^2 = -\chi^{-\frac{1}{2}}(r) \left(1 - \frac{r_0}{r}\right) dt^2 + \chi^{\frac{1}{2}}(r) \left[\left(1 - \frac{r_0}{r}\right)^{-1} dr^2 + \chi^{-\frac{1}{2}}(r) r^2 (d\theta^2 + \sin^2 \theta d\phi^2)\right], \quad (9)$$

where

$$\chi(r) = \left(1 - \frac{r_1}{r}\right)^{-2}. \quad (10)$$

The event horizon is at $r = r_0$ and we get the metrics

$$g_{\theta\theta} \rightarrow r^2 \left(1 - \frac{r_1}{r}\right) \quad \text{and} \quad g_{\phi\phi} \rightarrow r^2 \sin^2 \theta \left(1 - \frac{r_1}{r}\right). \quad (11)$$

So, the area of the black hole comes out to be **(author?)** [5, 6]

$$A = \int \sqrt{g_{\theta\theta} g_{\phi\phi}}|_{r=r_0} d\theta d\phi = 8\pi r_0 \sqrt{M^2 - Q^2}. \quad (12)$$

The horizon with visible area is possible, i.e., $A > 0$, only if $M > Q$. On the other hand, for an extremal black hole, $M = Q$ which corresponds to vanishing of the area, i.e., $A = 0$. So, the relation between entropy and area of a black hole given by equation(4) can never be obtained. This failure of the metric of equation(6) to reproduce the correct Bekenstein-Hawking entropy-area relationship(4) has prompted various attempts using D-branes**(author?)** [6] to reach out from higher dimensions to four and the concept of extended horizon**(author?)** [8]. Such attempts have been partially successful.

It is worth mentioning that in the metric components g_{tt} and g_{rr} in equation(5), the term $\frac{2M}{r}$ arises from gravitation and the term $\frac{Q^2}{r^2}$ arises from electromagnetic interaction, both of which are at a weaker scale. It is natural to expect an additional contribution arising from strong interaction effects that would modify the Reissner- Nordstrom metric and give a term $\sim \frac{1}{r^3}$. Indeed this leads to the correct entropy-area relation for an extremal black hole.

In section-2 the metric tensor for a spherical symmetric gravitational field is obtained. The Reissner-Nordstrom metric for a charged black hole is obtained in section-3, where the effect of electromagnetic interaction is taken into account. The section-4 is devoted to the modification of the Reissner-Nordstrom metric by introduction of an additional term arising from strong interaction Regge physics. In section-5 the area of the extremal black hole is obtained using the modified metric. The entropy of the black hole is calculated in section-6 and section-7 is devoted to concluding remarks.

2. THE METRIC TENSOR

As is well known, the metric tensor $g_{\mu\nu}(x)$ is related to the line element $d\tau$ by

$$d\tau^2 = g_{\mu\nu}(x) dx^\mu dx^\nu. \quad (13)$$

We give the details of the symmetry of the local geometrical object $g_{\mu\nu}(x)$ which is a symmetric covariant tensor. Let $g_{\mu\nu}(x)$ and $g'_{\mu\nu}(x)$ be the original and the transformed components of $g_{\mu\nu}(x)$ at the same point x of the spacetime manifold. For infinitesimal mappings of the manifold mapping group(MMG) (author?) [9], the symmetry implies

$$g'_{\mu\nu}(x) - g_{\mu\nu}(x) = \bar{\delta} g_{\mu\nu} \equiv -g_{\mu\rho} \xi^\rho_{,\nu} - g_{\rho\nu} \xi^\rho_{,\mu} - g_{\mu\nu,\rho} \xi^\rho = 0, \quad (14)$$

where $\xi^\rho_{,\nu} = \frac{\partial \xi^\rho}{\partial x^\nu}$. The four vectors ξ^ρ are the descriptors of the mapping and are the Killing vectors of the metric $g_{\mu\nu}(x)$. If $g_{\mu\nu} = \eta_{\mu\nu}$ at all points of the manifold, where $\eta_{\mu\nu}$ is the Minkowski metric, equation(14) becomes

$$\eta_{\mu\rho} \xi^\rho_{,\nu} + \eta_{\rho\nu} \xi^\rho_{,\mu} = 0. \quad (15)$$

The general solution of equation(15) can be written as

$$\xi^\rho = \varepsilon^\rho + b^\rho_\sigma x^\sigma. \quad (16)$$

Here ε^ρ and $b_{\rho\sigma} \equiv \eta_{\rho\mu} b^\mu_{\sigma} = -b_{\sigma\rho}$ are infinitesimal parameters. So, the symmetry group of the metric tensor $g_{\mu\nu}$ is the Poincare group.

The most general form of a spherically symmetric metric tensor $g_{\mu\nu}(x)$ is (author?) [9]

$$g_{00}(x) = f_1(r, t), \quad g_{0k}(x) = f_2(r, t) \quad \text{and} \quad g_{jk}(x) = f_3(r, t) \delta_{jk} + f_4(r, t) \frac{x^j x^k}{r^2}; \quad j, k = 1, 2, 3 \quad (17)$$

where f_1, f_2, f_3 and f_4 are arbitrary functions of r and t . In a finite region of spacetime it is always possible to reduce the most general spherically symmetric $g_{\mu\nu}$ to a form (author?) [9]

$$g_{00}(x) = a(r, t), \quad g_{0k}(x) = 0 \quad \text{and} \quad g_{jk}(x) = -\delta_{jk} + b(r, t) \frac{x^j x^k}{r^2}, \quad (18)$$

where $a(r, t)$ and $b(r, t)$ are functions of r and t .

With a point mass M located at the origin $r = 0$ and with $x^\mu = \{t, r, \theta, \phi\}$ the spherically symmetric, static space-time can be represented by the metric of the general form (author?) [10]

$$d\tau^2 = -e^{2\nu} dt^2 + e^{2\mu} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \quad (19)$$

where ν and μ are functions of r and t .

The non-vanishing tetrad components of Riemann tensor are

$$R_{1212} = R_{3131} = -\frac{1}{r} e^{-2\mu} \frac{\partial \mu}{\partial r}, \quad (20)$$

$$R_{2020} = R_{3030} = -\frac{1}{r} e^{-2\mu} \frac{\partial \nu}{\partial r}, \quad (21)$$

$$R_{1220} = R_{3030} = -\frac{1}{r} e^{-2\mu} \frac{\partial \nu}{\partial r}, \quad (22)$$

$$R_{3232} = -\frac{1}{r^2} (1 - e^{2\mu}), \quad (23)$$

and

$$R_{1010} = -e^{-\mu-\nu} \left[\frac{\partial}{\partial r} \left(e^{\nu-\mu} \frac{\partial \nu}{\partial r} \right) - \frac{\partial}{\partial t} \left(e^{-\nu+\mu} \frac{\partial \mu}{\partial t} \right) \right]. \quad (24)$$

The field equations follow by setting the various components of Ricci tensor $R_{\mu\nu}$ equal to zero. The equation

$$-R_{01} = R_{2021} + R_{3031} = \frac{2}{r} e^{-\nu-\mu} \frac{\partial \mu}{\partial t} = 0, \quad (25)$$

implies that μ is independent of t and is a function of only r , i.e., $\mu = \mu(r)$. Further, the equation

$$R_{00} + R_{11} = -R_{3030} - R_{2020} - R_{3131} - R_{1212} = 0, \quad (26)$$

gives

$$\frac{2}{r} e^{-2\mu} \frac{\partial (\mu + \nu)}{\partial r} = 0, \quad (27)$$

so that

$$\nu = -\mu(r) + f(t), \quad (28)$$

where $f(t)$ is independent of r and is an arbitrary function of time. By redefining time so that $e^{f(t)} dt$ is replaced by dt one can set $f(t)$ equal to zero and hence

$$\nu = -\mu(r). \quad (29)$$

Thus both ν and μ are independent of time and are functions of only r .

Further, the Einstein's equation

$$-R_{22} = R_{0202} + R_{1212} + R_{3232} = 0, \quad (30)$$

gives

$$2 \frac{e^{2\nu}}{r} \frac{\partial \nu}{\partial r} = \frac{1}{r^2} (1 - e^{2\nu}),$$

or,

$$(1 - e^{2\nu}) = 2r e^{2\nu} \frac{\partial \nu}{\partial r} = r \frac{\partial e^{2\nu}}{\partial r},$$

or,

$$d(r e^{2\nu}) = dr,$$

or,

$$r e^{2\nu} = r + \text{const} = r - 2M,$$

so that

$$e^{2\nu} = 1 - \frac{2M}{r} = e^{-2\mu}. \quad (31)$$

Here, the constant M is the mass located at the origin $r = 0$. So, the metric is

$$g_{tt} = -\left(1 - \frac{2M}{r}\right), \quad g_{rr} = \left(1 - \frac{2M}{r}\right)^{-1}, \quad g_{\theta\theta} = r^2 \quad \text{and} \quad g_{\phi\phi} = r^2 \sin^2 \theta. \quad (32)$$

Thus, in a spherically symmetric, static universe equation(32) represents the metric tensor $g_{\mu\nu}$ for a black hole of mass M . This is the Schwarzschild metric.

3. CONTRIBUTION FROM ELECTROMAGNETIC INTERACTION

If the black hole is electrically charged, the metric $g_{\mu\nu}$ is modified due to inclusion of extra term involving the charge of the black hole. In the presence of electromagnetic field, the Einstein's field equation of general relativity gets modified. The Ricci tensor $R_{\mu\nu}$ is no more equal to zero and has an electromagnetic source term,

$$R_{\mu\nu} = -2 \left(\eta^{\alpha\beta} F_{\mu\alpha} F_{\nu\beta} - \frac{1}{4} \eta_{\mu\nu} F_{\alpha\beta} F^{\alpha\beta} \right), \quad (33)$$

where $F_{\mu\alpha}$ is Maxwell's electromagnetic field tensor and $\eta_{\mu\nu}$ is the Minkowski metric. Because of spherical symmetry, the only non-zero component of the field strength tensor $F_{\mu\alpha}$ is $F_{01} = -\frac{Q}{r^2}$. So, in the Einstein's equation, instead of $R_{22} = 0$, we get $R_{22} = (F_{01})^2 = \frac{Q^2}{r^4}$ when electromagnetic interaction is taken into account. Now,

$$R_{22} = \frac{1}{r^2} (1 - e^{2\nu}) - 2 \frac{e^{2\nu}}{r} \frac{\partial \nu}{\partial r},$$

or,

$$r^2 R_{22} = (1 - e^{2\nu}) - r \frac{de^{2\nu}}{dr} = 1 - \frac{d}{dr} (r e^{2\nu}),$$

or,

$$d(r e^{2\nu}) = (1 - r^2 R_{22}) dr,$$

or,

$$r e^{2\nu} = r - \int_{\infty}^r r'^2 R_{22}(r') dr' + \text{const.}$$

or,

$$e^{2\nu} = 1 - \frac{2M}{r} - \frac{1}{r} \int_{\infty}^r r'^2 R_{22}(r') dr'. \quad (34)$$

where the constant is $-2M$. Using the value $R_{22}(r) = \frac{Q^2}{r^4}$, we get

$$e^{2\nu} = 1 - \frac{2M}{r} - \frac{1}{r} \int_{\infty}^r r'^2 \frac{Q^2}{r'^4} dr' = 1 - \frac{2M}{r} + \frac{Q^2}{r^2}. \quad (35)$$

Thus, for a static, spherically symmetric charge distribution, with total charge Q , the solution of Einstein's field equation leads to the Reissner-Nordstrom metric(**author?**) [10]

$$d\tau^2 = - \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right)^{-1} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2). \quad (36)$$

Unfortunately, this metric does not give the correct entropy-area relation for a charged, extremal black hole. The obvious reason is that the effect of strong interaction has not been taken into account, which would contribute a $\frac{1}{r^3}$ term to the metric coefficients as shown below.

4. CONTRIBUTION FROM THE STRONG INTERACTION

In order to obtain the correct entropy-area relation for a charged, extremal black hole, we modify the Reissner-Nordstrom metric of equation(36) by introducing an additional term arising from strong interaction. From strong interaction Regge physics we know that for a black hole of mass M^*

$$\alpha' M^{*2} = N, \quad (37)$$

where $N = 0, 1, 2, \dots$ and $\alpha' = \frac{1}{2}$ is the Regge slope. Here it is assumed that the mass $M = M^*$ of the black hole falls on the Regge trajectory or is very close to it.

In order to incorporate the effect of strong interaction, we consider the QCD Lagrangian(**author?**) [11]

$$L = -\frac{1}{4} F_{\alpha}^{\mu\nu} F_{\alpha\mu\nu} - \sum_n \bar{\psi}_n (\not{\partial} - ig \not{A}_{\alpha} t_{\alpha} + m_n) \psi_n, \quad (38)$$

where $F_{\alpha}^{\mu\nu}$ is the colour gauge covariant field strength tensor, A_{α}^{μ} is colour gauge vector potential, t_{α} are the generators of the colour $SU(3)$ group with $Tr(t_{\alpha} t_{\beta}) = \frac{1}{2} \delta_{\alpha\beta}$, n is the label for the quark flavour and g is the strong coupling constant. In order to investigate further, we note that a constant, background non-abelian gauge field $A_{\mu\nu}(x)$ cannot be removed by a gauge transformation.

For such a constant, static, background field, the field strength is(**author?**) [11]

$$F_{\alpha\mu\nu} = C_{\alpha\beta\gamma} A_{\beta\mu} A_{\gamma\nu}, \quad (39)$$

which has the gauge covariant derivative $D_{\lambda} F_{\delta\mu\nu} = C_{\delta\epsilon\gamma} C_{\alpha\beta\gamma} A_{\epsilon\lambda} A_{\beta\mu} A_{\lambda\nu}$. Here $C_{\alpha\beta\gamma}$ are the structure constants. The contribution of the constant, static background field can be inferred qualitatively from perturbation expansion so that we have a term with $\frac{1}{r^3}$ times the usual one with field $A_{\alpha\beta} A_{\beta\gamma} A_{\gamma\mu}$ having the coefficient

$$C_{\alpha\beta\gamma} C_{\beta\gamma\mu} \rightarrow (const) \delta_{\alpha\mu} \quad (40)$$

The non-linear terms arising from strong interaction give rise to additional contribution to the Reissner-Nordstrom metric as given below.

In the presence of both electromagnetic and strong interaction, the source term gets modified and the Ricci tensor R_{22} is now given by

$$R_{22} = \frac{Q^2}{r^4} - 2\lambda_v \frac{M^{*2}}{r^5} = \frac{Q^2}{r^4} - 2\lambda_v \frac{N}{\alpha' r^5}, \quad (41)$$

where M^* is the mass of the black hole and $\alpha' M^{*2} = N$ from equation(37).

The solution given in equation(34) now becomes

$$e^{2\nu} = 1 - \frac{2M}{r} - \frac{1}{r} \int_{\infty}^r r'^2 R_{22}(r') dr' = 1 - \frac{2M}{r} - \frac{1}{r} \int_{\infty}^r r'^2 \left(\frac{Q^2}{r^4} - 2\lambda_v \frac{N}{\alpha' r^5} \right) dr',$$

or,

$$e^{2\nu} = 1 - \frac{2M}{r} + \frac{Q^2}{r^2} - \lambda_v \frac{N}{\alpha'} \frac{1}{r^3}. \quad (42)$$

The Reissner-Nordstrom metric of equation(36) now gets an extra $\frac{1}{r^3}$ term and the modified metric is given by

$$d\tau^2 = - \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} - \lambda_v \frac{N}{\alpha'} \frac{1}{r^3} \right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} - \lambda_v \frac{N}{\alpha'} \frac{1}{r^3} \right)^{-1} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \quad (43)$$

The constant λ_v in equation(43) is determined from the condition that the black hole is to be extremal.

5. AREA OF THE BLACK HOLE

In order to find the area of the event horizon of the black hole, we need the metric coefficients $g_{\theta\theta}$ and $g_{\phi\phi}$ which, in turn, are obtained from the equation(43) by recasting it as follows.

We consider the cubic equation

$$1 - \frac{2M}{r} + \frac{Q^2}{r^2} - \lambda_v \frac{N}{\alpha'} \frac{1}{r^3} = 0, \quad (44)$$

whose roots r_0 , r_1 and r_2 satisfy the relations

$$r_0 + r_1 + r_2 = 2M, \quad r_1 r_2 + r_0 r_1 + r_0 r_2 = Q^2 \quad \text{and} \quad r_0 r_1 r_2 = \lambda_v \frac{N}{\alpha'}. \quad (45)$$

Equation(45) has the solutions

$$r_0 = \frac{2}{3}M - \frac{1}{6}\sqrt{4M^2 - 3Q^2 - \frac{3}{2}\frac{N}{\alpha'}} + \frac{1}{2}\sqrt{4M^2 - 3Q^2 + \frac{1}{2}\frac{N}{\alpha'}}, \quad (46)$$

$$r_1 = \frac{2}{3}M + \frac{1}{3}\sqrt{4M^2 - 3Q^2 - \frac{3}{2}\frac{N}{\alpha'}}, \quad (47)$$

and

$$r_2 = \frac{2}{3}M - \frac{1}{6}\sqrt{4M^2 - 3Q^2 - \frac{3}{2}\frac{N}{\alpha'}} - \frac{1}{2}\sqrt{4M^2 - 3Q^2 + \frac{1}{2}\frac{N}{\alpha'}}. \quad (48)$$

The solutions(46,47,48) will be real only if $4M^2 \geq (3Q^2 + \frac{3}{2}\frac{N}{\alpha'})$. In order to calculate the event horizon, we must have at least

$$4M^2 = \left(3Q^2 + \frac{3}{2}\frac{N}{\alpha'} \right). \quad (49)$$

For an extremal black hole, $M = Q$, and the condition given in equation(49) becomes

$$Q^2 = \frac{3}{2}\frac{N}{\alpha'}. \quad (50)$$

In this case, it is important to note that the area A of the extremal black hole will no more be equal to zero.

For an extremal black hole, the solutions (46), (47) and (48) become

$$r_0 = \frac{2}{3}M + \sqrt{\frac{N}{2\alpha'}}, \quad r_1 = \frac{2}{3}M \quad \text{and} \quad r_2 = \frac{2}{3}M - \sqrt{\frac{N}{2\alpha'}}. \quad (51)$$

Substituting these in the relation $r_0 r_1 r_2 = \lambda_v \frac{N}{\alpha'}$ of equation(45), we get

$$\lambda_v = \frac{M}{9}. \quad (52)$$

The metric of equation(43) now becomes

$$d\tau^2 = - \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} - \frac{MN}{9\alpha'} \frac{1}{r^3} \right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} - \frac{MN}{9\alpha'} \frac{1}{r^3} \right)^{-1} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2). \quad (53)$$

In terms of the solutions r_0 , r_1 and r_2 the metric can be recast as

$$d\tau^2 = -\chi^{-\frac{1}{2}}(r) \left(1 - \frac{r_0}{r} \right) dt^2 + \chi^{\frac{1}{2}}(r) \left[\left(1 - \frac{r_0}{r} \right)^{-1} dr^2 + \chi^{-\frac{1}{2}}(r) r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \right] \quad (54)$$

where $\chi^{-\frac{1}{2}}(r) = \left(1 - \frac{r_1}{r} \right) \left(1 - \frac{r_2}{r} \right)$. So, the metric coefficients are

$$g_{\theta\theta} \rightarrow r^2 \left(1 - \frac{r_1}{r} \right) \left(1 - \frac{r_2}{r} \right), \quad g_{\phi\phi} \rightarrow r^2 \left(1 - \frac{r_1}{r} \right) \left(1 - \frac{r_2}{r} \right) \sin^2 \theta. \quad (55)$$

From equation(51) we get the product

$$(r_0 - r_1)(r_0 - r_2) = \frac{N}{\alpha'} = 2N, \quad (56)$$

which is needed to calculate the area of the black hole. Here, $\alpha' = \frac{1}{2}$ is the Regge slope. The area of the event horizon of the black hole comes out to be

$$A = \int \sqrt{g_{\theta\theta} g_{\phi\phi}}|_{r=r_0} d\theta d\phi = 4\pi (r_0 - r_1)(r_0 - r_2) = 4\pi \frac{N}{\alpha'} = 4\pi M \sqrt{\frac{N}{\alpha'}} = 4\pi M \sqrt{2N}. \quad (57)$$

since $\frac{N}{\alpha'} = M^2$.

It may be noted that there is no multiplying factor like $(M^2 - Q^2)$ (as in equation(12))which tends to zero for an extremal black hole.

For a closed string like object, there will be two equations like equation(44) with the last terms being $\lambda_{v,L} \frac{N}{\alpha'} \frac{1}{r^3}$ and $\lambda_{v,R} \frac{N}{\alpha'} \frac{1}{r^3}$. So, $\chi^{-\frac{1}{2}}(r)$ will become

$$\chi_{L,R}^{-\frac{1}{2}}(r) = \left(1 - \frac{r_1}{r} \right) \left(1 - \frac{r_{2L,R}}{r} \right), \quad (58)$$

where $r_0 - r_{2L} = \frac{2N_L}{\alpha'} = 4N_L$ and $r_0 - r_{2R} = \frac{2N_R}{\alpha'} = 4N_R$. The area of the black hole becomes

$$A^{\text{close}} = 4\pi M \left(\sqrt{4N_L} + \sqrt{4N_R} \right) = 8\pi M \left(\sqrt{N_L} + \sqrt{N_R} \right). \quad (59)$$

6. ENTROPY OF THE BLACK HOLE

In order to calculate the entropy of the black hole we need the enumeration of the physical modes and have to take a 26 dimensional theory. In 26-D Nambu-Goto[12, 13] bosonic string, there are 24 physical bosons. Since the normal ordering constant for each boson is equal to $-\frac{1}{24}$, the total normal ordering constant has the value $a = -1$. The total number of open string bosonic states d_n , is described by the generating function

$$G(\omega) = \sum_{n=0}^{\infty} d_n \omega^n = \text{tr } \omega^N. \quad (60)$$

The generating function is evaluated by using the elementary methods of quantum statistical mechanics, namely **(author?)** [14]

$$\text{tr } \omega^N = \prod_{n=1}^{\infty} (1 - \omega^n)^{-N} = [f(\omega)]^{-N} = [f(\omega)]^{-24}, \quad (61)$$

where

$$f(\omega) = \prod_{n=1}^{\infty} (1 - \omega^n), \quad (62)$$

is the classical partition function. For estimation of the asymptotic density of states, the behaviour of the function $f(\omega)$, as $\omega \rightarrow 1$, is to be known. This is achieved by using

$$\begin{aligned} f(\omega) &= \exp \left(\sum_{n=1}^{\infty} \ln (1 - \omega^n) \right) = \exp \left(- \sum_{m,n=1}^{\infty} \frac{\omega^{mn}}{m} \right) = \exp \left(\sum_{m=1}^{\infty} \frac{\omega^m}{m(1 - \omega^m)} \right) \\ &\sim \exp \left(- \left(\frac{1}{1 - \omega} \right) \left(\sum_{m=1}^{\infty} \frac{1}{m^2} \right) \right) \sim \exp \left(- \left(\frac{1}{1 - \omega} \right) \frac{\pi^2}{6} \right). \end{aligned} \quad (63)$$

We can project out d_n from $G(\omega) = \sum_{n=0}^{\infty} d_n \omega^n$ by a contour integral along a small circle about the origin,

$$d_n = \frac{1}{2\pi i} \oint \frac{G(\omega)}{\omega^{n+1}} d\omega. \quad (64)$$

For large n , we have a sharply defined saddle point for ω near $\omega = 1$. One finds that **(author?)** [14], as $n \rightarrow \infty$

$$d_N \sim (\text{const.}) e^{\pi\sqrt{2N}}. \quad (65)$$

So, entropy of the extremal black hole, for open string, is

$$S = M \ln d_N = M \pi \sqrt{2N}. \quad (66)$$

From equations(66) and (57), the relation between entropy and area of the extremal black hole, for open string, comes out to be

$$S = \frac{A}{4}. \quad (67)$$

This result was obtained for $D = 26$, where we have used 26 bosonic coordinates in our theory. We now proceed to evaluate the entropy of a black hole using a four dimensional theory, as we should. For this, we consider a 4-D bosonic superstring constructed by one of us (BBD)**(author?)** [15]. Here the four bosonic coordinates are added to the remaining twenty two, i.e., $4 \times 11 = 44$ Majorana fermions that transform as Lorentz vectors - a bosonic representation of $SO(3,1)$. This is an exact substitution in conformity with Mandelstam's proof of the equivalence between fermions and bosons in the anomaly free 26 -D Nambu-Goto string**(author?)** [12, 13] and is an ingenious way of introducing fermionic matter to the Nambu-Goto bosonic string.

In order to have supersymmetry, one should add both conformal and superconformal ghosts to the neutrino-like Majorana term of the theory**(author?)** [16]. Let the left handed and right handed neutrinos be described by the bare mass term in the most general gauge invariant Lagrangian as

$$- \mathcal{L}_{bare} = \frac{1}{2} \sum_{\ell\ell'} B_{\ell\ell'} \bar{N}_{\ell L} N_{\ell' R} + h.c. \quad (68)$$

Now, using the identity $\bar{\nu}_{\ell L} N_{\ell' R} = \bar{\tilde{N}}_{\ell' L} \hat{\nu}_{\ell R}$, one gets

$$-\mathcal{L}_{mass} = \frac{1}{2} \left(\bar{\nu}_L \tilde{N}_L \right) \begin{pmatrix} 0 & M \\ M^T & B \end{pmatrix} \begin{pmatrix} \hat{\nu}_R \\ N_R \end{pmatrix} + h.c. \quad (69)$$

Here, M and B are $\mathcal{N} \times \mathcal{N}$ matrices for $\mathcal{N}$ generations of fermions, and ν_L , N_R etc. are $\mathcal{N}$ element column vectors containing fields from any generation. The mass matrix is symmetric and upon diagonalisation of the mass terms, one gets $2\mathcal{N}$ Majorana neutrinos. This can be seen by considering the simple case of one generation, i.e., $\mathcal{N} = 1$. Here, the mass matrix is $\mathcal{M} = \begin{pmatrix} 0 & M \\ M & B \end{pmatrix}$, where M and B are simple numbers. Using the orthogonal matrix $\mathcal{O} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ and the relation $\tan 2\theta = \frac{2M}{B}$, one can have

$$\mathcal{O} \mathcal{M} \mathcal{O}^T = \begin{pmatrix} -m_1 & 0 \\ 0 & m_2 \end{pmatrix} = \begin{pmatrix} m_1 & 0 \\ 0 & m_2 \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} = m \mathcal{K}^2. \quad (70)$$

Here m is the matrix with positive eigenvalues and $\mathcal{K}^2$ is a diagonal matrix of negative and positive signs. One finds that $\mathcal{M} = \mathcal{O}^T m \mathcal{K}^2 \mathcal{O}$. Now using the column vectors $\begin{pmatrix} n_{1L} \\ n_{2L} \end{pmatrix} = \mathcal{O} \begin{pmatrix} \nu_L \\ \hat{N}_L \end{pmatrix}$ and $\begin{pmatrix} n_{1R} \\ n_{2R} \end{pmatrix} = \mathcal{K}^2 \mathcal{O} \begin{pmatrix} \nu_L \\ \hat{N}_L \end{pmatrix}$, equation(69) becomes

$$-\mathcal{L}_{mass} = m_1 \bar{n}_{1L} n_{1R} + m_2 \bar{n}_{2L} n_{2R} + h.c. \quad (71)$$

Thus, the two eigenvalues obtained for a single generation are

$$n_1 = n_{1L} + n_{1R} = \cos \theta (\nu_L - \hat{\nu}_R) - \sin \theta (\hat{N}_L - N_R), \quad (72)$$

and

$$n_2 = n_{2L} + n_{2R} = \sin \theta (\nu_L + \hat{\nu}_R) + \cos \theta (\hat{N}_L + N_R). \quad (73)$$

Here $\hat{\nu}_R$ and $\hat{N}_L$ are conjugates of ν_L and N_R respectively. We thus get

$$n_1 = -\hat{n}_1 \quad \text{and} \quad n_2 = \hat{n}_2. \quad (74)$$

Both n_1 and n_2 are Majorana particles with $n_1 + n_2 = 11$ and $n_1 - n_2 = 1$.

The action which we consider is (author?) [16]

$$S = -\frac{1}{2\pi} \int d^2\sigma \left[\partial^\alpha X^\mu(\sigma, \tau) \partial_\alpha X_\mu(\sigma, \tau) - i \sum_{j=1}^6 \bar{\psi}^{\mu,j}(\sigma, \tau) \rho^\alpha \partial_\alpha \psi_{\mu,j}(\sigma, \tau) + i \sum_{k=1}^5 \bar{\phi}^{\mu,k} \bar{\psi}^{\mu,j}(\sigma, \tau) \rho^\alpha \partial_\alpha \phi_{\mu,k}(\sigma, \tau) \right]. \quad (75)$$

Here, X^μ is the bosonic field of the 26-D theory. The fermionic fields are ψ^μ and ϕ^μ , with $\partial_\alpha = (\partial_0, \partial_1) = (\partial_\sigma, \partial_\tau)$ and $\rho^0 \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$, $\rho^1 \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$; $\bar{\phi} = \phi^\dagger \rho^0$.

The value of the normal ordering constant $a = -1$, must remain unchanged in a superstring which is conformally and superconformally invariant. The number of bosons is to be calculated in such a way as to have the value of the normal ordering constant as $a = -1$. Out of the 44 fermionic modes, only 40 correspond to physical modes, because in all, there are six constraint equations - two constraints due to the vanishing of the energy momentum tensor, and four more constraints due to vanishing of the two species of currents due to the ψ and ϕ fermions in the action of equation(75). The 40 fermions account for $-\frac{40}{48}$ of the normal ordering constant and the remaining $-\frac{8}{48} = -4 \times \frac{1}{24}$ must be accounted for by four massless photon-like bosons. There are only two transverse coordinates, contributing $-\frac{4}{48}$. The longitudinal components which have been fermionised must be rebosonised. The reason is

that the Majorana fermions ϕ 's of the action(75) in the last term having the phase freedom for creation operators satisfying the anticommutation relation

$$\{\phi_\mu(\sigma, \tau), \phi_\nu^\dagger(\sigma', \tau)\} = -\eta_{\mu\nu} \delta(\sigma - \sigma'), \quad (76)$$

has a negative sign instead of the usual positive sign and ghosts in the normal sense. These fermions account for two current constraints out of the total four. It is possible to bosonise the fermions which are propagating on a circle(**author?**) [13]. One of the two rebosonised bosons $\phi^\dagger(\sigma)$ has the normal mode expansion for open strings

$$\phi^\dagger(\sigma) = \phi_0 + \sigma p_0 + i \sum_{n \neq 0} \frac{1}{n} \phi_n e^{in\sigma}, \quad (77)$$

and the Hamiltonian for one boson is

$$\mathcal{H} = \frac{p_0^2}{2} + \sum_n n \phi_{-n} \phi_n - \frac{1}{24}, \quad (78)$$

with $[p_0, \phi_0] = -i$ and $[\phi_n, \phi_m] = n\delta_{m+n}$, just like the transverse bosonic modes of the string with normal ordering for one boson as $-\frac{1}{24}$. Hence these must be counted as they are like the physical states of the larger of the two Fock spaces of the superconformal fields of the string theory in all aspects(**author?**) [17]. The normal ordering constant from these two rebosonised ones will be $-2 \times \frac{1}{24} = -4 \times \frac{1}{48}$, so that the value of the total comes out to $a = -1$.

The number of the physical fermionic level density is now

$$N_F^{\text{phys}} = \sum_{\mu=1,2} b_{-\frac{1}{2}}^{6,\mu} b_{\frac{1}{2},6,\mu} + \sum_{\mu=0,3} b_{-\frac{1}{2}}^{5,\mu} b'_{\frac{1}{2},5,\mu} + \sum_{\mu=0}^3 \sum_{j=1}^5 b_{-\frac{1}{2}}^{j,\mu} b_{\frac{1}{2},j,\mu} + \sum_{\mu=0}^3 \sum_{k=1}^4 b_{-\frac{1}{2}}^{k,\mu} b'_{\frac{1}{2},k,\mu} = 2 - 2 + 20 - 16 = 4, \quad (79)$$

since $b'_{-r} = -b_r^\dagger$ and $b_{-r} = b_r^\dagger$ are the quanta of the fermionic fields ψ and ϕ . Furthermore, the number of bosonic level density is

$$N_B^{\text{phys}} = N_B^{\text{tr}} + N_B^{\text{BFB}} = 2 + 2 = 4. \quad (80)$$

Here N_B^{tr} are the two transverse modes and N_B^{BFB} are the modes for the longitudinal bosons which had been fermionised and again rebosonised to account for the normal ordering constant for the entire enlarged Fock space. The above equality of both N_F^{phys} and N_B^{phys} to 4, thus ensures the supersymmetry of the physical states.

The degeneracy d_n is now obtained from the generating function

$$G(\omega) = \sum_n d_n \omega^n = \text{tr} \quad \omega^N = 4 \prod_{N=1}^{\infty} \left(\frac{1 + \omega^N}{1 - \omega^N} \right)^4, \quad (81)$$

with $\omega = e^{-2i\pi/r}$. Here, as explained earlier, $N_B = N_F = 4$ and the ground state degeneracy is also equal to 4. Asymptotically, i.e., as $\omega \rightarrow 1$, we have

$$G(\omega) \sim e^{2\pi^2/(1-\omega)}, \quad (82)$$

which yields

$$d_N = \frac{1}{2\pi i} \oint \frac{G(\omega)}{\omega^N + 1} d\omega \sim e^{\pi\sqrt{2N}}. \quad (83)$$

For a closed string, we have $\alpha' M^2 = 2(N_L + N_R)$ instead of $\alpha' M^2 = N$. In this case, the level densities, again being statistical,

$$d_N^{\text{close}} = d_{N_L} \cdot d_{N_R} \sim \exp \left(2\pi \left(\sqrt{N_L} + \sqrt{N_R} \right) \right). \quad (84)$$

The corresponding entropy is

$$S^{\text{close}} = M \ln d_N^{\text{close}} = 2\pi M \left(\sqrt{N_L} + \sqrt{N_R} \right). \quad (85)$$

From equations(85) and (59), we get the entropy-area relation for extremal black hole, for closed string, as

$$S^{\text{close}} = \frac{A^{\text{close}}}{4}. \quad (86)$$

Thus equations(67) and (86) reproduce the exact Bekenstein-Hawking relation between entropy and area of black holes.

7. CONCLUSIONS

It is interesting to note that the modified metric for the black hole gets contribution of the $\frac{1}{r}$ term from gravitation, the $\frac{1}{r^2}$ term from electromagnetism and the additional $\frac{1}{r^3}$ term from strong interaction. Thus all the known interactions are encompassed by the black hole physics, specially in obtaining the Bekenstein-Hawking entropy-area law.

It is worth mentioning that, the previous attempts to obtain the area of extremal black holes in four dimensions having failed, one has to consider D-branes and extended solitons. The black holes are treated in higher dimensions rather than in four dimensions. Our present result, in four dimensions, dispenses with the need to go over to five dimensions and extended solitons. The correct entropy-area relation comes from the known concepts of general relativity and high energy physics.

-
- [1] J. M. Bardeen, B. Carter and S. W. Hawking, Comm. Math. Phys.**31**(2),161 (1973), S. W. Hawking, Comm. Math. Phys.**43**(3), 199 (1975)
 - [2] J. D. Bekenstein, Lett. Nuov. Cim. **4**, 737 (1972), Phys. Rev. **D7**, 2333 (1973), Phys. Rev. **D9**, 3292 (1974)
 - [3] H.Reissner, Ann. Physik. Leipzig **50**, 106 (1916)
 - [4] G. Nordstrom, Proc. Con. Ned. Acad. Wet.**20**, 1238 (1918)
 - [5] R.H.Boyer and R.W. Lindquist, J. Math. Phys.**8**, 265 (1967)
 - [6] B.Carter, Phys. Rev. **174**, 1559 (1968)
 - [7] A. Strominger and C. Vafa, Phys. Lett.**B379**, 99 (1996), J. Maldacena and A. Strominger, hep-th/9603060, C.Johnson, R.Khuri and R.Myers, hep-th/9603061
 - [8] A. Sen, Mod. Phys. Lett. **A 10**, 2081 (1995)
 - [9] J. L. Anderson, *Principles of Relativity Physics*, (Academic Press, New York, 1967)
 - [10] S.Chandrasekhar, *The Mathematical Theory of Black holes*, (Oxford University Press, 1983)
 - [11] S. Weinberg, *The Quantum Theory of Fields, Vol-II*, (Cambridge University Press, New York, 1995)
 - [12] Y. Nambu, Lectures at Copenhagen Symposium, in *Symmetries and Quark models*, ed. R. Chand (Gordon and Breach, 1970)
 - [13] T. Goto, Prog. Theor. Phys.**46**, 1560 (1971)
 - [14] M. Green, J. H. Schwarz and E. Witten, *Superstring Theory, Vol-1* (Cambridge University Press, Cambridge, 1987)
 - [15] B. B. Deo, Phys. Lett. **B557**, 115 (2003)
 - [16] B. B. Deo, Mod. Phys. Lett.**A 21**, 6575 (2006); See also e.g., R. N. Mohapatra and P.B. Pal, *Massive neutrinos in Physics and Astrophysics*,(World Scientific, 1991)
 - [17] M. Kaku, *Introduction to Superstring Theory and M Theory*, (Springer, 1998)